VSOP AO2 PROPOSAL COVER SHEETS

DEADLINE: 8 May, 1998

SEND TO: VSOG, ISAS, 3-1-1 Yoshinodai, Sagamihara, Kanagawa 229-8510, JAPAN

Please read Appendix C of Announcement of Opportunity for details on how to fill in this Cover Sheet.

(1) Date prepared: 1998, April 24

(2) Proposal title: High Linear Resolution Imaging of z < 0.06 Sources

| (3) | INVESTIGATORS | INSTITUTION |
|-------|---|--|
| P.I. | S. Tingay | Jet Propulsion Laboratory, USA |
| co-I. | J. Reynolds, D. Jauncey, A. Tzioumis | Autsralia Telescope National Facility, Australia |
| co-I. | R. Preston, D. Murphy, D. Jones, D. Meier | Jet Propulsion Laboratory, USA |
| co-I. | P. McCulloch, D. Moffett, M. Costa | University of Tasmania, Australia |
| co-I. | G. Nicolson | Hartebeestoek RAO, South Africa |
| co-I. | | |

(4) Principal Investigator (or contact person) details...

Name: Steven J. Tingay

Address: Jet Propulsion Laboratory

E-mail: tingay@hyaa.jpl.nasa.gov : MS238-332

Fax : +1 818 393 6890 : 4800 Oak Grove Drive Phone : +1 818 354 4909 : Pasadena, CA 91109

: USA

(5) Proposal Abstract:

We have compiled a sample of the closest extragalactic objects in the Southern hemisphere ($|b| > 10^{\circ}$) within a redshift of z < 0.06, and containing compact radio components suitable for imaging with VSOP.

VSOP imaging of nearby active galaxies at 5GHz will return the highest linear resolution possible with the mission.

The list comprises a diversity of source types, including GHz-Peaked-Spectrum (GPS), FR-I, FR-II, BL Lac and γ -ray sources. Observations with VSOP at 5GHz will give linear resolution of 0.3pc or better on all objects in the sample - matching or improving existing observations of these objects in any waveband. We propose repeat observations at an interval of approximately six months to detect any structural evolution.

| (6) Proposal Category (indicate all that apply): Object type: ✓ AGN, ☐ Maser, ☐ Stellar, ☐ Pulsar, ☐ Other: Observation type: ✓ Continuum, ☐ Spectral Line, ☐ Polarization, ☐ Time-critical, ☐ Other: | these objects in any waveband. We propose repeat observations at an interval of approximately six months to detect any structural evolution. |
|--|--|
| | Object type: V AGN, Maser, Stellar, Pulsar, Other: Observation type: |

(7) Number of proposed experiments

An 'experiment' is one or more observations of one source at a fixed HALCA set-up. A request to observe the same source at 1.6 GHz and separately at 5 GHz requires two columns to be filled in in item (8) below. A request to observe the same source with HALCA simultaneously observing at 1.6 GHz and 5 GHz requires one column to be filled in. Multi-epoch observations of the same source at the same frequency – a 'monitoring experiment' – requires only one column to be filled in. Suggested observing dates, especially for for time-critical and monitoring experiments, should be specified in item (10).

The number of experiments in this proposal is: 6

(8) Details of proposed experiments

| | Experiment 1 | Experiment 2 | Experiment 3 | Experiment 4 |
|---------------------------------------|---------------|-------------------|---------------|---------------|
| Source name $(Jhhmm \pm ddmm)$ | J0519-4546 | J0522-3627 | J1325-4301 | J1517-2422 |
| Alternative name | Pictor A | | Centaurus A | AP Libræ |
| RA(J2000) (hh mm ss.ssss) | 05 19 49.75 | 05 22 57.985 | 13 25 27.620 | 15 17 41.814 |
| Dec(J2000) (dd mm ss.ssss) | -45 46 43.800 | -36 27 30.850 | -43 01 08.800 | -24 22 19.473 |
| Observing frequency band (GHz) | 5 | 5 | 5 | 5 |
| Continuum observations: | | | | |
| Standard VSOP freq. channels? | | | | |
| Channel A range (MHz) | | | | |
| Channel B range (MHz) | | | | |
| Spectral line observations: | | | | |
| Ch.A spectral line rest freq. (MHz) | | | | |
| Ch.A LSR velocity (km/s) | | | | |
| Ch.B spectral line rest freq. (MHz) | | | | |
| Ch.B LSR velocity (km/s) | | | | |
| FWHM of field of view required (mas) | | | | |
| Min. spectral channels per IF channel | 128 | 128 | 128 | 128 |
| Correlator averaging time (sec) | | | | |
| No. of correlating passes $(if > 1)$ | | | | |
| Total flux density (Jy) | 15.5 | 9.3 | 681 | 3.0 |
| Correlated flux (mJy) | 1100 | 1200 | 5500 | 2200 |
| Ground Radio Telescopes: | | | | |
| Suggested array given at Item (10)? | | | | |
| GRT observing mode: | | | | |
| 128Mbps LCP (standard) | | | | |
| $128 \mathrm{Mbps}\ \mathrm{LCP/RCP}$ | | | | |
| 256Mbps LCP/RCP | | | | |
| Preferred correlator: | | | | |
| No preference | | | | |
| Mitaka | | | | |
| Penticton | | | | |
| Socorro | | $ \overline{} $ | | $ ar{ abla} $ |
| Monitoring programs: | | | | |
| Number of observations | 2 | 2 | 2 | 2 |
| Mean interval (days) | ~200 | ~200 | ~200 | ~200 |
| Related AO1 proposal code(s) | V112 | V112 | | |

| | Experiment 5 | Experiment 6 | Experiment 7 | Experiment 8 |
|---------------------------------------|---------------|--------------|--------------|--------------|
| Source name $(Jhhmm \pm ddmm)$ | J1723-6500 | J2157-6941 | | |
| Alternative name | NGC 6328 | | | |
| RA(J2000) (hh mm ss.ssss) | 17 23 41.029 | 21 57 06.129 | | |
| Dec(J2000) (dd mm ss.ssss) | -65 00 36.610 | -69 41 24.52 | | |
| Observing frequency band (GHz) | 5 | 5 | | |
| Continuum observations: | | | | |
| Standard VSOP freq. channels? | | | | |
| Channel A range (MHz) | | | | |
| Channel B range (MHz) | | | | |
| Spectral line observations: | | | | |
| Ch.A spectral line rest freq. (MHz) | | | | |
| Ch.A LSR velocity (km/s) | | | | |
| Ch.B spectral line rest freq. (MHz) | | | | |
| Ch.B LSR velocity (km/s) | | | | |
| FWHM of field of view required (mas) | | | | |
| Min. spectral channels per IF channel | 128 | 128 | | |
| Correlator averaging time (sec) | | | | |
| No. of correlating passes $(if > 1)$ | | | | |
| Total flux density (Jy) | 3.8 | 12.3 | | |
| Correlated flux (mJy) | 3800 | 1200 | | |
| Ground Radio Telescopes: | | | | |
| Suggested array given at Item (10)? | $ \nabla$ | | | |
| GRT observing mode: | | | | |
| 128Mbps LCP (standard) | | | | |
| $128 \mathrm{Mbps}\ \mathrm{LCP/RCP}$ | | | | |
| $256 \mathrm{Mbps}\ \mathrm{LCP/RCP}$ | | | | |
| Preferred correlator: | | | | |
| No preference | | | | |
| Mitaka | | | | |
| Penticton | | | | |
| Socorro | | | | |
| Monitoring programs: | | | | |
| Number of observations | 2 | 2 | | |
| Mean interval (days) | ~200 | ~200 | | |
| Related AO1 proposal code(s) | V101 | V112 | | |

| (9) VSOP spacecraft observing mode (see Section 3 and Table 5 of the VSOP Proposer's Guide): |
|--|
| $\boxed{\hspace{-0.1cm} \bigvee}$ 2 channel x 16 MHz, 2-bit (Standard mode), Other: |
| Phase calibration tones: |
| ∇ On (Standard continuum mode), |
| Off (Standard spectral line mode) |
| (Include justification of any non-standard choice at (10) below) |
| |
| (10) Additional notes to the scheduler: |
| We request an array of Southern Hemisphere telescopes consisting of the ATCA, Mopra, Hobart, Hartebeestoek, and Ceduna for those sources south of $\delta = -45^{\circ}$. For all other sources, we request a combination of two arrays, the VLBA and the Southern Hemisphere array given above. For these combined experiments, if the volume of tape copying required for processing at a single correlator is too great, we request that the separate components of the experiments be correlated on separate correlators. |
| |
| |
| |
| (11) Attach a scientific and technical justification, not in excess of 2 pages of text and 2 pages of figures. Up to one page of (u,v) plots per source may optionally be included. (Refer to the VSOP Announcement of Opportunity for detailed instructions.) Preprints and reprints will not be forwarded to the Scientific Review Committee. |
| Send two paper copies of the complete proposal to: |
| VSOP Observing Proposals |
| VSOP Science Operations Group |

Information from the Cover Sheets of scheduled proposals will be made available from the VSOP WWW site.

Proposals must be received at ISAS by 8 May 1998

3-1-1 Yoshinodai, Sagamihara Kanagawa 229-8510 JAPAN

Institute of Space and Astronautical Science

In addition, e-mail the completed LATEX file to submit@vsop.isas.ac.jp